

Page 1/2

# Heard and McDonald Islands linthorst energy

Island and McDonald Islands on the Central Kerguelen Plateau. Flares were observed with a split-beam ...

????????(?:Heard Island and McDonald Islands,?:HIMI),????????,1947????????,1997????????(????????)??????1600????????;????????1000????????,????3700?????,??????80%????????,?????? ...

Heard Island and McDonald Islands (HIMI) are a sub-Antarctic island group. They are located in the Southern Ocean, about 4,000 km south west of mainland Australia. Penguins are by far the most abundant birds on the island.

????????(?: Heard Island and McDonald Islands,?:HIMI),????????,1947????? ???? ,1997????????(?????? ...

The activity on the McDonald Islands group resulted in the island doubling in size and increasing in elevation by about 100 m (Australian Government 2016). Along with the French Kerguelen group of islands 440 km to the northwest, Heard and McDonald Islands form the only exposed parts of the submarine Kerguelen plume (Australian Government 2016).

Island and McDonald Islands group show unique evolutionary adaptations to the environment, and several other invertebrate groups provide opportunities to study evolutionary processes in undisturbed populations at the southern limits

????????(?: Heard Island and McDonald Islands,?:HIMI),????????,1947????? ???? ,1997????????(??????)?

Web: <https://gennergyps.co.za>